

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.


Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.


**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

FIG. 1



X=Cl Radical (1)

X=H Monocillin I (2)



Geldanamycin (3)

FIG.2





FIG.3

3/28



- (a) TBDPSCI, imid., >95%; (b) DIBAL-H, -78 °C, 92%;
 (c) LiCl, DIPEA (EtO)₂P(O)CH₂CO₂Et, 95%;
 (d) DIBAL-H, -20 °C, 96%; (e) (+)-DET, Ti(OiPr)₄, TBHP, 90%, >95% ee; (f) SO₃*pyridine, Et₃N, DMSO, 90%;
 (g) PH₃PCH₃Br, NaHMDS, 0 °C, 82%; (h) TBAF, 89%.




FIG.4




FIG.5







FIG. 6



a. *n*-BuLi, -78 °C, 50% (6:1); b. TBSCl, 83%; c. 42 °C, 70%; d. (i) mCPBA, (ii) Ac₂O, Et₃N, H₂O, 60 °C, (iii) NaHCO₃, MeOH, 60%; e. SO₂Cl₂, 50%

c



FIG. 8




FIG. 9

9/28









FIG.10



a. TBSCl, pyridine; b. NIS or NBS, TsOH; c. Pd(PPh)₃, RSnBu₃, d. nBu₄NF


FIG. 11-1



TO FIG. 11-2

FROM FIG. 11-1

FIG. 11-2




7

39

44

12/28



22

45

46






FIG. 12-1




TO FIG. 12-2

FROM FIG. 12-1


FIG. 12-2



1. Rainey Ni





51




52

14/28




53



54


FIG. 13

15/28




^a (a) TBDPSCl, imid., >95%; (b) DIBAL-H, -78 °C, 92%; (c) LiCl, DIPEA (EtO)₂P(O)CH₂CO₂Et, 95%; (d) DIBAL-H -20 °C, 96%; (e) (+)-tetramethyltartaric acid diamide-BBu, Et₂Zn, CH₂I₂, 9 >95% ee; (f) SO₃*pyridine, Et₃N, DMSO, 90%; (g) Ph₃PCH NaHMDS, 0 °C, 82%; (h) TBAF, 89%; (i) 7, P(furyl)₃, DIA benzene, 60%

FIG. 14



a. n-BuLi, -78 °C, 75% (3:1); b. TBSCl, 83%; c. 42 °C, 20%; d. (i) mCPBA, (ii) Ac₂O, Et₃N, H₂O, 60 °C, (iii) NaHCO₃, MeOH, 60%; e. SO₂Cl₂, 80%



FIG. 15-1




TO FIG. 15-2

FROM FIG. 15-1

FIG. 15-2




18/28



GD=Geldanamycin




FIG. 16-1




TO FIG. 16-2

FROM FIG. 16-1

FIG. 16-2



Radicalol, Steroid, GD




Radicalol, Steroid, GD


FIG. 17-1

21/28


I. Radical




III. Cyclopropyl radical



II. Monocillin I




IV. Cyclopropyl monocillin



TO FIG. 17-2

FIG. 17-2

MCF7 Cells Treated with Radicicol and Analogues




HER2


FROM FIG. 17-2

FIG. 17-3


I. Radicol



VII. Radicol Oxime



V. Dimethyl Monocillin I



VI. Dimethyl Radicol






FIG. 18-1

24/28


I. Radicol




III. Cyclopropyl radicol



II. Monocillin I



IV. Cyclopropyl monocillin




TO FIG. 18-2

FROM FIG. 18-1

FIG. 18-2

BT474 Cells Treated with Novel Radicicoliols (24hrs.)

25/28



HER2

FIG.19

Growth of MCF7 Treated with Radicicol and Derivatives of Radicicol




FIG.20





FIG. 21



Therapeutic Effect of Cycloproparadicol in Nude Mice Bearing Human Mammary Carcinoma MX-1 Xenograft (Q2Dx7, iv.injection)




FIG. 22




FIG. 23

MX-1 tumors
12 hrs following a 6 hr CIVI





FIG. 24

IC₅₀ of Growth Inhibition of Different Tumor Cell lines




Cell lines

MCF-7 54 nM
 CCRF-CEM 42 nM
 CCRF-CEM/Vinblastin resistant 32 nM

>500 nM

>500 nM





Cell lines

MCF-7 150 nM, >500 nM
 CCRF-CEM 208 nM
 CCRF-CEM/Vinblastin resistant 209 nM

390 nM

>500 nM
 >5 μM



Cell lines
 MCF-7

98 nM

282 nM




FIG. 25

Degradation of HER2 by Cycloproparadicicol Analogues




FIG. 26




FIG. 27